

THIS CONTENT IS PROTECTED AND MAY NOT BE SHARED, UPLOADED, OR DISTRIBUTED.
FOR USE ONLY OF UC MERCED CSE176 SPRING 2023 STUDENTS.

Exercise 1: kernel machines (18 points). Consider the XOR binary classification problem, with a training set $\{(\mathbf{x}_n, y_n)\}_{n=1}^N \subset \mathbb{R}^2 \times \{-1, +1\}$ given by $\left\{ \left(\begin{pmatrix} -1 \\ +1 \end{pmatrix}, +1 \right), \left(\begin{pmatrix} +1 \\ -1 \end{pmatrix}, +1 \right), \left(\begin{pmatrix} -1 \\ -1 \end{pmatrix}, -1 \right), \left(\begin{pmatrix} +1 \\ +1 \end{pmatrix}, -1 \right) \right\}$, which is not linearly separable. Construct a SVM to learn a nonlinear discriminant function as follows.

- (2 points) Define a feature function $\mathbf{z} = \phi(\mathbf{x}) = \begin{pmatrix} x_1 x_2 \\ x_2 \end{pmatrix} \in \mathbb{R}^2$. Evaluate it at each training point to obtain $\mathbf{z}_n = \phi(\mathbf{x}_n)$ and plot them in the new space \mathbf{z} . Verify the points are now linearly separable.
- (8 points) Find a linear discriminant $G(\mathbf{z}) = \mathbf{w}^T \mathbf{z} + w_0$ such that $\text{sgn}(G(\phi(\mathbf{x}_n))) = y_n \forall n$. Make sure this discriminant has the maximum margin, give the value of the margin, find the support vectors, and plot the separating hyperplane and the SVs. *Hint:* there is no need to solve any QP, just use geometric intuition. Explain your answer.
- (6 points) Write the nonlinear discriminant $g(\mathbf{x}) = \mathbf{w}^T \phi(\mathbf{x}) + w_0$ in the original \mathbf{x} -space. Draw in the plane: the boundary $g(\mathbf{x}) = 0$, and the curves corresponding to each of the training points (i.e., the sets $\{\mathbf{x} \in \mathbb{R}^2: g(\mathbf{x}) = g(\mathbf{x}_n)\}$ for each $n = 1, \dots, 4$).
- (2 points) Relate this result to the solution of the primal and dual QP for a kernel SVM. Specifically, write the kernel function $K(\mathbf{x}, \mathbf{y})$ that results from the choice of ϕ above, the general form of the kernel SVM discriminant function $g(\mathbf{x})$, and guess what the values of the Lagrange multipliers α_n should be if we solved the dual QP. *Hint:* again, use geometric intuition and symmetry.

Exercise 2: kernel machines (18 points). Consider binary classification for $\mathbf{x} \in \mathbb{R}^2$ using the nonlinear discriminant function $g(\mathbf{x}) = x_1^2 + x_2^2 - 1$ and assigning a label $y = \text{sgn}(g(\mathbf{x})) \in \{-1, +1\}$ to an instance \mathbf{x} .

- (4 points) g splits the plane into two class regions. Draw them (or explain them in plain English). Are they linearly separable?

Consider a new feature function $\phi: \mathbf{x} \in \mathbb{R}^2 \rightarrow \mathbb{R}^D$, which creates D features made up (in some way) of the original features x_1, x_2 .

- (10 points) Define a function ϕ such that, using the new features $\mathbf{z} = \phi(\mathbf{x}) \in \mathbb{R}^D$, we can define a linear discriminant function $G(\mathbf{z}) = \mathbf{w}^T \mathbf{z}$ that is equivalent to g , i.e., $G(\phi(\mathbf{x})) = g(\mathbf{x})$. Relate this to kernel functions and support vector machines.
- (4 points) Define what the kernel function $K(\cdot, \cdot)$ would be that corresponds to your feature function ϕ .

Exercise 3: graphical models (6 points). Consider the following two graphical models defined on binary random variables $X, Y, Z \in \{0, 1\}$, given by their joint distributions:

$$p(X, Y, Z) = p(X)p(Z|X,Y)p(Y|X) \quad \text{and} \quad p(X, Y, Z) = p(X)p(Z)p(Y|Z)$$

For each of them:

- (4 points) Prove that $\sum_{X,Y,Z} p(X, Y, Z) = 1$.
- (2 points) Draw the graphical model.

Exercise 4: graphical models (21 points).

Consider 3 binary random variables with joint distribution given by the table.

- (14 points) Evaluate the following probabilities (notation: X means $X = 1$, \bar{X} means $X = 0$): $p(X)$, $p(Z)$, $p(X, Z)$, $p(X|Y)$, $p(Z|Y)$, $p(Y|X)$, and $p(X, Z|Y)$.
- (4 points) Show by direct evaluation that this distribution has the property that X and Z are marginally dependent, i.e., $P(X, Z) \neq p(X)p(Z)$ (for all values of X and Z); but that they become independent when conditioned on Y , i.e., $p(X, Z|Y) = p(X|Y)p(Z|Y)$ for all values of X, Z and Y .
- (3 points) Show by direct evaluation that $p(X, Z, Y) = p(X)p(Y|X)p(Z|Y)$ (for all values of X and Z). Draw the corresponding directed graph for this graphical model.


X	Y	Z	$p(X, Y, Z)$
0	0	0	0.192
0	0	1	0.048
0	1	0	0.144
0	1	1	0.216
1	0	0	0.192
1	0	1	0.048
1	1	0	0.064
1	1	1	0.096

Show your work in all cases.

Exercise 5: graphical models (21 points).

Consider a graphical model defined on binary random variables (where variables X_i correspond to diseases and variables Y_j to symptoms), given by the following diagram and conditional probability tables at each node.

Note: in the tables and the questions, the notation “ $p(Y_3|\bar{X}_1, X_2)$ ” means “ $p(Y_3 = 1|X_1 = 0, X_2 = 1)$ ”, etc.




conditional probability tables at each node				
X_1 (“flu”)	X_2 (“hayfever”)	Y_1 (“fever”)	Y_2 (“headache”)	Y_3 (“fatigue”)
$p(X_1) = 0.4$	$p(X_2) = 0.1$	$p(Y_1 X_1) = 0.8$	$p(Y_2 X_1, X_2) = 0.9$	$p(Y_3 X_1, X_2) = 0.7$
		$p(Y_1 \bar{X}_1) = 0.1$	$p(Y_2 X_1, \bar{X}_2) = 0.8$	$p(Y_3 X_1, \bar{X}_2) = 0.7$
			$p(Y_2 \bar{X}_1, X_2) = 0.7$	$p(Y_3 \bar{X}_1, X_2) = 0.3$
			$p(Y_2 \bar{X}_1, \bar{X}_2) = 0.1$	$p(Y_3 \bar{X}_1, \bar{X}_2) = 0.7$

- (3 points) Give the expression of the joint distribution it defines over all the variables.
- (18 points) Calculate the value of the following probabilities:
 - $p(\bar{Y}_2|X_1, \bar{X}_2)$.
 - $p(Y_1, Y_3|\bar{X}_1, \bar{X}_2)$.
 - $p(Y_1|X_2)$.
 - $p(Y_1)$.
 - $p(X_1|Y_1, \bar{Y}_2)$.
 - $p(X_2|\bar{Y}_1, Y_2, \bar{Y}_3)$.

Show your work in all cases.

Exercise 6: discrete Markov models (11 points).

Consider the discrete Markov model given by the diagram.



- (3 points) Give the set of states of this discrete Markov model, its transition matrix \mathbf{A} and its vector of initial state probabilities $\boldsymbol{\pi}$.
- (8 points) Compute the probability of the following sequences: 1123, 313321 and 3213.

Show your work in all cases.

Exercise 7: discrete Markov models (7 points). Consider a discrete Markov model with two states a, b .

- (5 points) We have a training set consisting of the following sequences: **aaaa, bbabb, ababa, bababa**. Give the maximum likelihood estimate of the parameters $(\mathbf{A}, \boldsymbol{\pi})$.
- (2 points) Draw the corresponding discrete Markov model as in the previous exercise.

Show your work in all cases.