

This set covers chapters 16–19 of the book *Numerical Optimization* by Nocedal and Wright, 2nd ed.

From the book, the following exercises: 16.1–3, 16.6, 16.10, 16.15–16, 16.20–21, 17.4, 17.11, 18.5, 19.12–13. In addition, the exercises below. There are no Matlab programming exercises. However, you may find it useful to plot the objective function, constraints and auxiliary functions (e.g. the quadratic-penalty function) with `fcontour`.

IV.1. Quadratic programming. Consider the Markowitz model of portfolio optimization (example 16.1 in the book):

$$\max_{\mathbf{x} \in \mathbb{R}^n} q(\mathbf{x}) = \mathbf{x}^T \boldsymbol{\mu} - \kappa \mathbf{x}^T \mathbf{G} \mathbf{x} \quad \text{subject to} \quad \mathbf{e}^T \mathbf{x} = 1, \mathbf{x} \geq \mathbf{0}$$

where $\kappa \geq 0$, $\boldsymbol{\mu} > \mathbf{0}$, \mathbf{G} is symmetric positive definite and $\mathbf{e} = (1, \dots, 1)^T$. W.l.o.g. assume that the largest component of $\boldsymbol{\mu}$ is the first ($\mu_1 \geq \mu_i, i = 1, \dots, n$).

- (i) Suppose that $g_{ii} > g_{ij}, i = 1, \dots, n$. Show that, for a solution to be at one corner of the feasible polytope, namely $\mathbf{x}^* = (1, 0, \dots, 0)^T$, the following condition must hold:

$$\kappa \leq \kappa_a \text{ with } \kappa_a = \min_{i=2, \dots, n} \frac{\mu_1 - \mu_i}{2(g_{11} - g_{1i})}.$$

Interpret this solution. (Hint: apply the KKT and 2nd-order conditions to \mathbf{x}^* .)

- (ii) Suppose that the sum of each column of \mathbf{G}^{-1} is positive. Show that, for a solution to have only positive components ($\mathbf{x} > \mathbf{0}$), the following condition must hold:

$$\kappa > \kappa_c \text{ with } \kappa_c = \frac{1}{2} \mathbf{e}^T \mathbf{G}^{-1} \boldsymbol{\mu} - \frac{1}{2} \mathbf{e}^T \mathbf{G}^{-1} \mathbf{e} \left(\min_{i=1, \dots, n} \frac{(\mathbf{G}^{-1} \boldsymbol{\mu})_i}{(\mathbf{G}^{-1} \mathbf{e})_i} \right).$$

What is the solution \mathbf{x}^* ? What is the solution for $\kappa \rightarrow \infty$?

- (iii) Consider a different portfolio optimization problem:

$$\min_{\mathbf{x} \in \mathbb{R}^n} \mathbf{x}^T \mathbf{G} \mathbf{x} \quad \text{subject to} \quad \boldsymbol{\mu}^T \mathbf{x} \geq \kappa, \mathbf{e}^T \mathbf{x} = 1, \mathbf{x} \geq \mathbf{0}$$

where $\kappa \geq 0$ as before. What is the largest κ for which this problem is feasible? (Hint: formulate as an LP $\kappa = \max \boldsymbol{\mu}^T \mathbf{x}$ s.t. $\mathbf{e}^T \mathbf{x} = 1, \mathbf{x} \geq \mathbf{0}$; guess its solution and prove it is a solution using the KKT conditions.)

IV.2. Quadratic-penalty, augmented-Lagrangian, log-barrier and interior-point methods.

- (i) Consider the constrained optimization problem $\min_{\mathbf{x}} x_1^2 + x_2^2$ s.t. $x_1 + x_2 = 1$. Find the solution $(\mathbf{x}^*, \lambda^*)$ of this problem using the KKT and second-order conditions. Write the quadratic-penalty function $Q(\mathbf{x}; \mu)$ and its gradient $\nabla_{\mathbf{x}} Q(\mathbf{x}; \mu)$ and Hessian $\nabla_{\mathbf{x}\mathbf{x}}^2 Q(\mathbf{x}; \mu)$. Show that: $Q(\mathbf{x}; \mu)$ has a single minimiser \mathbf{x}_k for each $\mu_k > 0$; this minimiser tends to the solution \mathbf{x}^* of the problem as $\mu_k \rightarrow \infty$; the Lagrange multiplier estimate $\lambda_k \approx -\mu_k c(\mathbf{x}_k)$ (eq. (17.10) in the book) tends to the Lagrange multiplier λ^* at the solution as $\mu_k \rightarrow \infty$; the Hessian of the penalty function at the minimiser becomes progressively more ill-conditioned as $\mu_k \rightarrow \infty$, i.e., $\text{cond}(\nabla_{\mathbf{x}\mathbf{x}}^2 Q(\mathbf{x}_k; \mu_k)) \rightarrow \infty$.
- (ii) As in (i) but using the augmented-Lagrangian function $\mathcal{L}_A(\mathbf{x}, \lambda; \mu)$ where the variable λ_k is updated as $\lambda_{k+1} \leftarrow \lambda_k - \mu_k c(\mathbf{x}_k)$ (eq. (17.39) in the book). Also, prove that the sequence of iterates $(\mathbf{x}_k, \lambda_k)$ converges to the solution even if μ_k is kept at a constant value $\mu > 0$ (i.e., we do not drive $\mu_k \rightarrow \infty$), and give the convergence order. How does it depend on μ ?
- (iii) As in (i) but for the problem $\min_{\mathbf{x}} x_1^2 + x_2^2$ s.t. $x_1 \geq 1$ using the log-barrier function $P(\mathbf{x}; \mu)$ and where the Lagrange multiplier estimate is $\lambda_k \approx \frac{\mu_k}{c(\mathbf{x}_k)}$ (eq. (19.47) in the book).
- (iv) For the problem in (iii), write the system of perturbed KKT equations $\mathbf{F}(\dots) = \mathbf{0}$ for an interior-point method; solve it and find the primal-dual central path; compute the Jacobian \mathbf{J} of \mathbf{F} and indicate how to obtain the Newton step; show that \mathbf{J} does not become progressively more ill-conditioned as we approach the solution.

IV.3. Convex equality-constrained quadratic programming using quadratic-penalty and augmented-Lagrangian methods. Consider the strictly convex equality-constrained QP:

$$\min_{\mathbf{x} \in \mathbb{R}^n} \frac{1}{2} \mathbf{x}^T \mathbf{G} \mathbf{x} + \mathbf{c}^T \mathbf{x} \quad \text{subject to} \quad \mathbf{A} \mathbf{x} = \mathbf{b}$$

where \mathbf{G} is symmetric pd and $\mathbf{A}_{m \times n}$ has full rank with $m < n$.

- (i) Solve the QP, giving an explicit expression for the solution, i.e., the KKT point $(\mathbf{x}^*, \boldsymbol{\lambda}^*)$.
- (ii) Write the quadratic-penalty function $Q(\mathbf{x}; \mu)$, prove it is strictly convex and find its minimizer $\mathbf{x}(\mu)$ and its Lagrange multiplier estimates $\boldsymbol{\lambda}(\mu) = -\mu(\mathbf{A}\mathbf{x}(\mu) - \mathbf{b})$ explicitly. Prove $\mathbf{x}(\mu) \xrightarrow{\mu \rightarrow \infty} \mathbf{x}^*$ and $\boldsymbol{\lambda}(\mu) \xrightarrow{\mu \rightarrow \infty} \boldsymbol{\lambda}^*$. Hint: use the Sherman-Morrison-Woodbury formula.
- (iii) Write the augmented-Lagrangian function $\mathcal{L}_A(\mathbf{x}, \boldsymbol{\lambda}; \mu)$, prove it is strictly convex and find its minimizer $\mathbf{x}(\boldsymbol{\lambda}; \mu)$ and the update for $\boldsymbol{\lambda}$ explicitly. Prove $\mathbf{x}(\boldsymbol{\lambda}; \mu) \xrightarrow{\mu \rightarrow \infty} \mathbf{x}^*$ and $\boldsymbol{\lambda}(\mu) \xrightarrow{\mu \rightarrow \infty} \boldsymbol{\lambda}^*$.

Now we keep the penalty parameter fixed to a value $\mu > 0$. Prove that iterating the update for $\boldsymbol{\lambda}$ also converges to the solution, and determine the convergence order. Hint: $\|\mathbf{M}\mathbf{u}\| \leq \|\mathbf{M}\| \|\mathbf{u}\|$, using the L_2 norm for vectors and matrices, i.e., $\|\mathbf{u}\| = \sqrt{u_1^2 + \dots + u_n^2}$ and $\|\mathbf{M}\| =$ largest singular value of \mathbf{M} .

IV.4. Generalized spectral problem. Prove the following theorem. Consider the optimisation problem

$$\max_{\mathbf{X}} \text{tr}(\mathbf{X} \mathbf{A} \mathbf{X}^T) \quad \text{s.t.} \quad \mathbf{X} \mathbf{B} \mathbf{X}^T = \mathbf{I} \quad (1)$$

where $\mathbf{X} \in \mathbb{R}^{L \times N}$, $L < N$, $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{N \times N}$, \mathbf{A} is symmetric and \mathbf{B} is symmetric positive definite. Let $\mathbf{C} = \mathbf{B}^{-\frac{1}{2}} \mathbf{A} \mathbf{B}^{-\frac{1}{2}}$ have spectral representation $\mathbf{C} = \mathbf{U} \boldsymbol{\Lambda} \mathbf{U}^T$ and assume its eigenvalues are distinct though not necessarily positive. Then, the solution of (1) is unique and given by $\mathbf{X} = \mathbf{U}_L^T \mathbf{B}^{-\frac{1}{2}}$, where $\mathbf{U}_L = (\mathbf{u}_1, \dots, \mathbf{u}_L)$ are the leading L eigenvectors of \mathbf{C} .